

# Kinto<sup>®</sup> Plus

**THIS SEED HAS BEEN TREATED WITH Kinto Plus (contains triticonazole, fluxapyroxad and fludioxonil) MAPP 19506.**

Sowing treated seed that has been stored for prolonged periods (beyond the season of treatment) may adversely affect effectiveness and/or crop safety.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALS) AND SUITABLE PROTECTIVE GLOVES when handling treated seed and contaminated seed sowing equipment.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

DO NOT HANDLE seed unnecessarily.

DO NOT APPLY TREATED SEED FROM THE AIR.

DO NOT USE TREATED SEED as food or feed.

DO NOT RE-USE SACKS OR CONTAINERS THAT HAVE BEEN USED FOR TREATED SEED for food or feed.

KEEP TREATED SEED SECURE from people, domestic stock/pets and wildlife at all times during storage and use.

WASH HANDS AND EXPOSED SKIN before meals and after work.

TO PROTECT BIRDS AND WILD MAMMALS, treated seed should not be left on the soil surface. Bury or remove spillages; ensure that the product is also fully incorporated at the end of rows.

Treated seed should not be broadcast.

Seed should be drilled to a depth of 40 mm into a well-prepared seed bed.

If seed is present on the soil surface, or if spills have occurred, then, if conditions are appropriate, the field should be harrowed and then rolled to ensure good incorporation.

BASF plc  
Agricultural Solutions  
4th and 5th Floors  
2 Stockport Exchange  
Railway Road, Stockport,  
SK1 3GG  
Telephone 0161 475 3000  
Emergency Information: 0049 1802 273112 (24 Hours freephone)  
Technical Enquiries: 0845 602 2553 (Office hours)

 **BASF**  
We create chemistry